

SEMISTABLE REDUCTION IN CHARACTERISTIC 0 FOR FAMILIES OF SURFACES AND THREE-FOLDS

K. KARU

1. INTRODUCTION

In [N-K] the semistable reduction of a morphism $F : X \rightarrow B$ was stated as a problem in the combinatorics of polyhedral complexes. In this paper we solve it in the case when the relative dimension of F is no bigger than three.

First we recall the setup of the problem from [N-K]. The ground field k will be algebraically closed of characteristic zero.

Definition 1.1. A flat morphism $F : X \rightarrow B$ of nonsingular projective varieties is semistable if in local analytic coordinates $x_1, \dots, x_n$ at $x \in X$ and $t_1, \dots, t_m$ at $b \in B$ the morphism F is given by

$$t_i = \prod_{j=l_{i-1}+1}^{l_i} x_j$$

where $0 = l_0 < l_1 < \dots < l_m \leq n$.

The conjecture of semistable reduction states that

Conjecture 1.2. Let $F : X \rightarrow B$ be a surjective morphism with geometrically integral generic fiber. There exist an alteration (proper surjective generically finite morphism) $B' \rightarrow B$ and a modification (proper birational morphism) $X' \rightarrow X \times_B B'$ such that $X' \rightarrow B'$ is semistable.

Conjecture 1.2 was proved in [KKMS] (main theorem of Chapter 2) in case when B is a curve. A weak version of the conjecture was proved in [N-K] for arbitrary X and B . In both cases the proof proceeds by reducing F to a morphism of toroidal embeddings, stating the problem in terms of the associated polyhedral complexes, and solving the combinatorial problem.

1.1. Polyhedral complexes. We consider (rational, conical) polyhedral complexes $\Delta = (|\Delta|, \{\sigma\}, \{N_\sigma\})$ consisting of a collection of lattices $N_\sigma \cong \mathbb{Z}^n$ and rational full cones $\sigma \subset N_\sigma \otimes \mathbb{R}$ with a vertex. The cones σ are glued together to form the space $|\Delta|$ so that the usual axioms of polyhedral complexes hold:

1. If $\sigma \in \Delta$ is a cone, then every face σ' of σ is also in Δ , and $N_{\sigma'} = N_\sigma|_{\text{Span}(\sigma')}$.
2. The intersection of two cones $\sigma_1 \cap \sigma_2$ is a face of both of them, $N_{\sigma_1 \cap \sigma_2} = N_{\sigma_1}|_{\text{Span}(\sigma_1 \cap \sigma_2)} = N_{\sigma_2}|_{\text{Span}(\sigma_1 \cap \sigma_2)}$.

A morphism $f : \Delta_X \rightarrow \Delta_B$ of polyhedral complexes $\Delta_X = (|\Delta_X|, \{\sigma\}, \{N_\sigma\})$ and $\Delta_B = (|\Delta_B|, \{\tau\}, \{N_\tau\})$ is a compatible collection of linear maps $f_\sigma : (\sigma, N_\sigma) \rightarrow (\tau, N_\tau)$; i.e. if σ' is a face of σ then $f_{\sigma'}$ is the restriction of f_σ . We will only consider morphisms $f : \Delta_X \rightarrow \Delta_B$ such that $f_\sigma^{-1}(0) \cap \sigma = \{0\}$ for all $\sigma \in \Delta_X$.

Definition 1.3. A surjective morphism $f : \Delta_X \rightarrow \Delta_B$ such that $f^{-1}(0) = \{0\}$ is semistable if

1. Δ_X and Δ_B are nonsingular.
2. For any cone $\sigma \in \Delta_X$, we have $f(\sigma) \in \Delta_B$ and $f(N_\sigma) = N_{f(\sigma)}$.

We say that f is weakly semistable if it satisfies the two properties except that Δ_X may be singular.

The following two operations are allowed on Δ_X and Δ_B :

1. Projective subdivisions Δ'_X of Δ_X and Δ'_B of Δ_B such that f induces a morphism $f' : \Delta'_X \rightarrow \Delta'_B$;
2. Lattice alterations: let $\Delta'_X = (|\Delta_X|, \{\sigma\}, \{N'_\sigma\})$, $\Delta'_B = (|\Delta_B|, \{\tau\}, \{N'_\tau\})$, for some compatible collection of sublattices $N'_\tau \subset N_\tau$, $N'_\sigma = f^{-1}(N'_\tau) \cap N_\sigma$, and let $f' : \Delta'_X \rightarrow \Delta'_B$ be the morphism induced by f .

Conjecture 1.4. Given a surjective morphism $f : \Delta_X \rightarrow \Delta_B$, such that $f^{-1}(0) = \{0\}$, there exists a projective subdivision $f' : \Delta'_X \rightarrow \Delta'_B$ followed by a lattice alteration $f'' : \Delta''_X \rightarrow \Delta''_B$ so that f'' is semistable.

$$\begin{array}{ccccc} \Delta_{X''} & \rightarrow & \Delta_{X'} & \rightarrow & \Delta_X \\ \downarrow f'' & & \downarrow f' & & \downarrow f \\ \Delta_{B''} & \rightarrow & \Delta_{B'} & \rightarrow & \Delta_B \end{array}$$

The importance of Conjecture 1.4 lies in the fact that it implies Conjecture 1.2 (Proposition 8.5 in [N-K]). In the case when $\dim(\Delta_B) = 1$, Conjecture 1.4 was proved in [KKMS] (main theorem of Chapter 3). In [N-K] (Theorem 0.3) the conjecture was proved with semistable replaced by weakly semistable. The main result of this paper is

Theorem 1.5. *Conjecture 1.4 is true if f has relative dimension ≤ 3 . Hence, Conjecture 1.2 is true if F has relative dimension ≤ 3 .*

The relative dimension of a linear map $f : \sigma \rightarrow \tau$ of cones σ, τ is $\dim(\sigma) - \dim(f(\sigma))$. The relative dimension of $f : \Delta_X \rightarrow \Delta_B$ is by definition the maximum of the relative dimensions of $f_\sigma : \sigma \rightarrow \tau$ over all $\sigma \in \Delta_X$. If $F : X \rightarrow B$ is a morphism of toroidal embeddings of relative dimension d , then the associated morphism of polyhedral complexes $f : \Delta_X \rightarrow \Delta_B$ has relative dimension $\leq d$ because in local models the relative dimension of F is no bigger than the rank of the kernel of $f : N_\sigma \rightarrow N_\tau$. Thus, the second statement of the theorem follows from the first.

1.2. Notation. We will use notations from [KKMS] and [F]. For a cone $\sigma \in N \otimes \mathbb{R}$ we write $\sigma = \langle v_1, \dots, v_n \rangle$ if $v_1, \dots, v_n$ lie on the 1-dimensional edges of σ and generate it. If v_i are the first lattice points along the edges we call them primitive points of σ . For a simplicial cone σ with primitive points $v_1, \dots, v_n$, the multiplicity of σ is

$$m(\sigma, N_\sigma) = [N_\sigma : \mathbb{Z}v_1 \oplus \dots \oplus \mathbb{Z}v_n].$$

A polyhedral complex Δ is nonsingular if and only if $m(\sigma, N_\sigma) = 1$ for all $\sigma \in \Delta$. To compute the multiplicity of σ we can count the representatives $w \in N_\sigma$ of classes of $N_\sigma / \mathbb{Z}v_1 \oplus \dots \oplus \mathbb{Z}v_n$ in the form

$$w = \sum_i \alpha_i v_i, \quad 0 \leq \alpha_i < 1.$$

Such points w were called Waterman points of σ in [KKMS]. Also notice that since the multiplicity of a face of σ is no bigger than the multiplicity of σ , to compute the multiplicity of Δ it suffices to consider maximal cones only.

If Δ_X and Δ_B are simplicial, we say that $f : \Delta_X \rightarrow \Delta_B$ is simplicial if $f(\sigma) \in \Delta_B$ for all $\sigma \in \Delta_X$. Assume that f is simplicial. Let $u_1, \dots, u_n$ be the primitive points of Δ_B , and $m_1, \dots, m_n$ positive integers. By taking the $(m_1, \dots, m_n)$ sublattice at $u_1, \dots, u_n$ we mean the lattice alteration $N'_\tau = \mathbb{Z}[m_{i_1}u_{i_1}, \dots, m_{i_l}u_{i_l}]$ where $\tau \in \Delta_B$ has primitive points $u_{i_1}, \dots, u_{i_l}$.

For cones $\sigma_1, \sigma_2 \in \Delta$ we write $\sigma_1 \leq \sigma_2$ if σ_1 is a face of σ_2 .

A subdivision Δ' of Δ is called projective if there exists a homogeneous piecewise linear function $\psi : |\Delta| \rightarrow \mathbb{R}$ taking rational values on the lattice points (a good function for short) such that the maximal cones of Δ' are exactly the maximal pieces in which ψ is linear.

1.3. Acknowledgment. The suggestion to write up the proof of semistable reduction for low relative dimensions came from Dan Abramovich.

2. JOINS

For cones $\sigma_1, \sigma_2 \in \mathbb{R}^N$ lying in complementary planes: $\text{Span}(\sigma_1) \cap \text{Span}(\sigma_2) = \{0\}$, the join of σ_1 and σ_2 is $\sigma_1 * \sigma_2 = \sigma_1 + \sigma_2$. Let σ be a simplicial cone $\sigma = \sigma_1 * \dots * \sigma_n$. If σ'_i is a subdivision of σ_i for all $i = 1, \dots, n$, we define the join

$$\sigma' = \sigma'_1 * \dots * \sigma'_n$$

as the set of cones $\rho = \rho_1 + \dots + \rho_n$, where $\rho_i \in \sigma'_i$.

Let $f : \Delta_X \rightarrow \Delta_B$ be a simplicial map of simplicial complexes. For u_i a primitive point of Δ_B , $i = 1, \dots, n$, let $\Delta_{X,i} = f^{-1}(\mathbb{R}_+ u_i)$ be the simplicial subcomplex of Δ_X . If $\Delta'_{X,i}$ is a subdivision of $\Delta_{X,i}$ for $i = 1, \dots, n$, we can define the join

$$\Delta'_X = \Delta'_{X,1} * \dots * \Delta'_{X,n}$$

by taking joins inside all cones $\sigma \in \Delta_X$. This is well defined by the assumption that $f^{-1}(0) = \{0\}$.

Lemma 2.1. *If $\Delta'_{X,i}$ are projective subdivisions of $\Delta_{X,i}$ then the join Δ'_X is a projective subdivision of Δ_X .*

Proof. Let ψ_i be good functions for $|\Delta'_{X,i}|$. Extend ψ_i linearly to the entire $|\Delta'_X|$ by setting $\psi_i(|\Delta'_{X,j}|) = 0$ for $j \neq i$. Clearly, $\psi = \sum_i \psi_i$ is a good function defining the subdivision Δ'_X . $\square$

Consider $f|_{\Delta_{X,i}} : \Delta_{X,i} \rightarrow \mathbb{R}_+ u_i$. By the main theorem of Chapter 2 in [KKMS] there exist a subdivision $\Delta'_{X,i}$ of $\Delta_{X,i}$ and an $m_i \in \mathbb{Z}$ such that after taking the m_i -sublattice at u_i we have $f'|_{\Delta'_{X,i}}$ semistable. Now let Δ'_X be the join of $\Delta'_{X,i}$, and take the $(m_1, \dots, m_n)$ -sublattice at $(u_1, \dots, u_n)$. Then $f' : \Delta'_X \rightarrow \Delta'_B$ is a simplicial map and $f'|_{\Delta'_{X,i}}$ is semistable.

We can also see that the multiplicity of Δ'_X is not bigger than the multiplicity of Δ_X . Let $\sigma \in \Delta_X$ have primitive points v_i and let $\sigma' \subset \sigma$ be a maximal cone in the subdivision with primitive points v'_i . The multiplicity of σ' is the number of Waterman points $w' \in N'_\sigma$

$$w' = \sum_i \alpha_i v'_i, \quad 0 \leq \alpha_i < 1.$$

We show that the set of Waterman points of σ' can be mapped injectively into the set of Waterman points of σ , hence the multiplicity of σ' is not bigger than the multiplicity of σ . Write

$$w' = \sum_i (\beta_i + b_i) v_i, \quad 0 \leq \beta_i < 1, \quad b_i \in \mathbb{Z}_+.$$

Then $w = \sum_i \beta_i v_i \in N_\sigma$ is a Waterman point of σ . If different w'_1, w'_2 give the same w , then $w'_1 - w'_2 \in N'_\sigma \cap \mathbb{Z}\{v_i\} = \mathbb{Z}\{v'_i\}$, hence $w'_1 - w'_2 = 0$.

3. MODIFIED BARYCENTRIC SUBDIVISIONS

Let $f : \Delta_X \rightarrow \Delta_B$ be a simplicial morphism of simplicial complexes. Consider the barycentric subdivision $BS(\Delta_B)$ of Δ_B . The 1-dimensional cones of $BS(\Delta_B)$ are $\mathbb{R}_+ \hat{\tau}$ where $\hat{\tau} = \sum u_i$ is the barycenter of a cone $\tau \in \Delta_B$ with primitive points $u_1, \dots, u_m$. A cone $\tau' \in BS(\Delta_B)$ is spanned by $\hat{\tau}_1, \dots, \hat{\tau}_k$, where $\tau_1 \leq \tau_2 \leq \dots \leq \tau_k$ is a chain of cones in Δ_B .

In general, f does not induce a morphism $BS(\Delta_X) \rightarrow BS(\Delta_B)$. For that we need to modify the barycenters $\hat{\sigma}$ of cones $\sigma \in \Delta_X$.

Definition 3.1. The data of **modified barycenters** consists of

1. A subset of cones $\tilde{\Delta}_X \subset \Delta_X$.
2. For each $\sigma \in \tilde{\Delta}_X$ a point $b_\sigma \in \text{int}(\sigma) \cap N_\sigma$ such that $f(b_\sigma) \in \mathbb{R}_+ \hat{\tau}$ for some $\tau \in \Delta_B$.

Recall that for any total order $\prec$ on the set of cones in Δ_X refining the partial order $\leq$, the barycentric subdivision $BS(\Delta_X)$ can be realized as a sequence of star subdivisions at the barycenters $\hat{\sigma}$ of $\sigma \in \Delta_X$ in the descending order $\prec$.

Definition 3.2. Given modified barycenters $(\tilde{\Delta}_X, \{b_\sigma\})$ and a total order $\prec$ on Δ_X refining the partial order $\leq$, the **modified barycentric subdivision** $MBS_{\tilde{\Delta}_X, \{b_\sigma\}, \prec}(\Delta_X)$ is the sequence of star subdivisions at b_σ for $\sigma \in \tilde{\Delta}_X$ in the descending order $\prec$.

To simplify notations, we will write $MBS(\Delta_X)$ instead of $MBS_{\tilde{\Delta}_X, \{b_\sigma\}, \prec}(\Delta_X)$. By definition, $MBS(\Delta_X)$ is a projective simplicial subdivision of Δ_X . As in the case of the ordinary barycentric subdivision, the cones of $MBS(\Delta_X)$ can be characterized by chains of cones in Δ_X . We may assume that the 1-dimensional cones of Δ_X are all in $\tilde{\Delta}_X$. For a cone $\sigma \in \Delta_X$ let $\tilde{\sigma}$ be the maximal face of σ (w.r.t. $\prec$) in $\tilde{\Delta}_X$. Given a chain of cones $\sigma_1 \leq \dots \leq \sigma_k$ in Δ_X , the cone spanned by $b_{\tilde{\sigma}_1}, \dots, b_{\tilde{\sigma}_k}$ is a subcone of σ_k . Let $C(\Delta_X)$ be the set of all such cones corresponding to chains $\sigma_1 \leq \dots \leq \sigma_k$ in Δ_X .

Proposition 3.3. $C(\Delta_X) = MBS(\Delta_X)$.

Proof. Let $BS(\Delta_X)$ be the ordinary barycentric subdivision of Δ_X . Both $C(\Delta_X)$ and $MBS(\Delta_X)$ are obtained from $BS(\Delta_X)$ by moving the barycenters $\hat{\sigma}$ (and everything attached to them) to the new position $b_{\tilde{\sigma}}$ for all $\sigma \in \Delta_X$ in the descending order $\prec$. $\square$

Corollary 3.4. If $f(\tilde{\sigma}) = f(\sigma)$ for all $\sigma \in \Delta_X$ then f induces a simplicial map $f' : MBS(\Delta_X) \rightarrow BS(\Delta_B)$.

Proof. Let $\sigma' \in MBS(\Delta_X)$ correspond to a chain $\sigma_1 \leq \dots \leq \sigma_k$. Then we have a chain of cones $f(\sigma_1) \leq \dots \leq f(\sigma_k)$ in Δ_B . The assumption that $f(\tilde{\sigma}_i) = f(\sigma_i)$ implies that $f(b_{\tilde{\sigma}_i}) \in \mathbb{R}_+ \hat{f}(\sigma_i)$, hence the cone $\langle b_{\tilde{\sigma}_1}, \dots, b_{\tilde{\sigma}_k} \rangle$ maps onto the cone $\langle \hat{f}(\sigma_1), \dots, \hat{f}(\sigma_k) \rangle \in BS(\Delta_B)$. $\square$

The hypothesis of the corollary is satisfied, for example, if for any $\sigma \in \Delta_X$ with $f(\sigma) = \tau \in \Delta_B$ and for any face $\sigma_1 \leq \sigma$ such that $\sigma_1 \in \tilde{\Delta}_X$, $f(\sigma_1) \neq \tau$, there exists $\sigma_2 \in \tilde{\Delta}_X$ such that $\sigma_1 \leq \sigma_2 \leq \sigma$ and $f(\sigma_2) = \tau$:

$$\begin{array}{ccccc} \sigma_1 & \leq & \sigma_2 & \leq & \sigma \\ \downarrow & & \downarrow & & \downarrow \\ \tau_1 & \leq & \tau & = & \tau \end{array}$$

Indeed, $\tilde{\sigma} \neq \sigma_1$ because $\sigma_1 \prec \sigma_2$.

3.1. Example. Assume that $f : \Delta_X \rightarrow \Delta_B$ is a simplicial map of simplicial complexes taking primitive points of Δ_X to primitive points of Δ_B (e.g. Δ_X is simplicial and f is weakly semistable). Then for a cone $\sigma \in \Delta_X$ such that $f : \sigma \xrightarrow{\sim} \tau$, we have $f(\hat{\sigma}) = \hat{\tau}$.

Let $\tilde{\Delta}_X = \bar{\Delta}_X = \{\sigma \in \Delta_X : f|_{\sigma} \text{ is injective}\}$, $b_{\sigma} = \hat{\sigma}$. In this case $\tilde{\sigma}$ is the maximal face of σ (w.r.t. $\prec$) such that $f|_{\sigma}$ is injective. Clearly, the hypothesis of the lemma is satisfied, and we have a simplicial map $f' : MBS(\Delta_X) \rightarrow BS(\Delta_B)$.

Next we compute the multiplicity of $MBS(\Delta_X)$. Let $\sigma \in \Delta_X$ have primitive points $v_1, \dots, v_n$, and let $\sigma' \subset \sigma$ be a maximal cone in the subdivision, corresponding to the chain

$$\langle v_1 \rangle \leq \langle v_1, v_2 \rangle \leq \dots \leq \langle v_1, \dots, v_n \rangle.$$

Since $\tilde{\rho} \subset \rho$ for any ρ , the primitive points of σ' can be written as

$$\begin{aligned} v'_1 &= a_{11}v_1 \\ v'_2 &= a_{21}v_1 + a_{22}v_2 \\ &\dots \\ v'_n &= a_{n1}v_1 + \dots + a_{nn}v_n \end{aligned}$$

for some $0 \leq a_{ij}$. The multiplicity of σ' is $a_{11} \cdot a_{22} \cdots a_{nn}$ times the multiplicity of σ . In case when b_{ρ} are barycenters $\hat{\rho}$, all $a_{ij} \leq 1$, hence the multiplicity of σ' is not bigger than the multiplicity of σ .

4. REDUCING THE MULTIPLICITY OF Δ_X

Let $f : \Delta_X \rightarrow \Delta_B$ be weakly semistable and Δ_X simplicial (i.e. Δ_B is nonsingular, Δ_X is simplicial, and f is a simplicial map taking primitive points of Δ_X to primitive points of Δ_B). Notice that if $\tilde{\Delta}_X$ is as in Example 3.1, then $\tilde{\Delta}_X$ is nonsingular, and $f(\hat{\sigma}) = \hat{f}(\sigma)$ for any $\sigma \in \tilde{\Delta}_X$.

A singular simplicial cone $\sigma \in \Delta_X$ with primitive points $v_1, \dots, v_n$ contains a Waterman point $w \in N_{\sigma}$,

$$w = \sum_i \alpha_i v_i, \quad 0 \leq \alpha_i < 1, \quad \sum_i \alpha_i > 0.$$

The star subdivision of σ at w has multiplicity strictly less than the multiplicity of σ .

We will show in this section that if every singular cone of Δ_X contains a Waterman point w mapping to a barycenter of Δ_B , then there exists a modified barycentric subdivision $MBS(\Delta_X)$ having multiplicity strictly less than the multiplicity of Δ_X , such that f induces a simplicial map $f' : MBS(\Delta_X) \rightarrow BS(\Delta_B)$.

For every singular cone $\sigma \in \Delta_X$ choose a point w_{σ} as follows. By assumption, there exists a Waterman point $w \in \sigma$ mapping to a barycenter of Δ_B : $f(w) = \hat{\tau}$. Write $f(\sigma) = \tau * \tau_0$ and choose a face $\sigma_0 \leq \sigma$ such that $f : \sigma_0 \xrightarrow{\sim} \tau_0$. Set $w_{\sigma} = w + \hat{\sigma}_0$; then

$$f(w_{\sigma}) = f(w) + f(\hat{\sigma}_0) = \hat{\tau} + \hat{\tau}_0 = \hat{f}(\sigma)$$

Having chosen the set $\{w_\sigma\}$, we may remove some of the points w_σ if necessary so that every simplex $\rho \in \Delta_X$ contains at most one w_σ in its interior. With $\bar{\Delta}_X$ as in Example 3.1, let $\tilde{\Delta}_X = \bar{\Delta}_X \cup \{\rho \in \Delta_X \mid w_\sigma \in \text{int}(\rho) \text{ for some singular } \sigma\}$, $b_\rho = \hat{\rho}$ if $\rho \in \bar{\Delta}_X$, and $b_\rho = w_\sigma$ if $w_\sigma \in \text{int}(\rho)$.

By construction, $(\tilde{\Delta}_X, \{b_\rho\})$ satisfies the hypothesis of Corollary 3.4, hence f induces a simplicial map $f' : MBS(\Delta_X) \rightarrow BS(\Delta_B)$. Before we compute the multiplicity of $MBS(\Delta_X)$, we choose a particular total order $\prec$ on Δ_X . Extend $\leq$ on Δ_X to a partial order $\prec_0$ by declaring that $\sigma_1 \prec_0 \sigma_2$ for all (nonsingular) $\sigma_1 \in \bar{\Delta}_X$ and singular $\sigma_2 \in \Delta_X$. Let $\prec$ be an extension of $\prec_0$ to a total order on Δ_X . With such $\prec$, if $\sigma \in \Delta_X$ is singular, then $b_{\tilde{\sigma}}$ is one of the points w_ρ .

As in Example 3.1, the multiplicity of $MBS(\Delta_X)$ is not bigger than the multiplicity of Δ_X . If $\sigma \in \Delta_X$ is singular we show by induction on the dimension of σ that the multiplicity of $MBS(\sigma)$ is strictly less than the multiplicity of σ . Let $v_1, \dots, v_N$ be the primitive points of σ , and consider the cone $\sigma' = \langle b_{\tilde{\sigma}}, v_1, \dots, v_{N-1} \rangle$ in the star subdivision of σ at $b_{\tilde{\sigma}} = \sum_i a_i v_i$. To show that every maximal cone of $MBS(\sigma)$ contained in σ' has multiplicity less than the multiplicity of σ , we have three cases:

1. If $a_N = 0$, then σ' is degenerate.
2. If $0 < a_N < 1$, then the multiplicity of $\langle b_{\tilde{\sigma}}, v_1, \dots, v_{N-1} \rangle$ is less than the multiplicity of σ , and since all $b_\rho = \sum_i c_i v_i$ have coefficients $0 \leq c_i \leq 1$, further subdivisions at b_ρ do not increase the multiplicity of $\langle b_{\tilde{\sigma}}, v_1, \dots, v_{N-1} \rangle$.
3. If $a_N = 1$, then $b_{\tilde{\sigma}} = w + \hat{\rho}$ for some $\rho \leq \sigma$ and $w \in \langle v_1, \dots, v_{N-1} \rangle$ a Waterman point. Hence $\langle v_1, \dots, v_{N-1} \rangle$ is singular and, by induction, every maximal cone in $MBS(\langle v_1, \dots, v_{N-1} \rangle)$ has multiplicity less than the multiplicity of $\langle v_1, \dots, v_{N-1} \rangle$. Then also every maximal cone in $\mathbb{R}_+ b_{\tilde{\sigma}} * MBS(\langle v_1, \dots, v_{N-1} \rangle)$ has multiplicity less than the multiplicity of $\langle b_{\tilde{\sigma}}, v_1, \dots, v_{N-1} \rangle$.

5. FAMILIES OF SURFACES AND 3-FOLDS.

Proof of Theorem 1.5. It is not difficult to subdivide Δ_X and Δ_B so that Δ_X is simplicial, Δ_B is nonsingular, and $f : \Delta_X \rightarrow \Delta_B$ is a simplicial map (e.g. Proposition 4.4 and the remark following it in [N-K]). Applying the join construction we can make $f|_{\Delta_{X,i}}$ semistable without increasing the multiplicity of Δ_X . We will show below that every singular simplex of Δ_X contains a Waterman point mapping to a barycenter of Δ_B . By the previous section, there exist a modified barycentric subdivision and a simplicial map $f' : MBS(\Delta_X) \rightarrow BS(\Delta_B)$, with multiplicity of $MBS(\Delta_X)$ strictly less than the multiplicity of Δ_X . Since f' is simplicial and $BS(\Delta_B)$ nonsingular, the proof is completed by induction.

Restrict f to a singular simplex $f : \sigma \rightarrow \tau$, where σ has primitive points $v_{ij}, i = 1, \dots, n, j = 1, \dots, J_i$, τ has primitive points $u_1, \dots, u_n$, and $f(v_{ij}) = u_i$. Since σ is singular, it contains a Waterman point

$$w = \sum_{i,j} \alpha_{ij} v_{ij}, \quad 0 \leq \alpha_{ij} < 1,$$

where not all $\alpha_{ij} = 0$. Restricting to a face of σ if necessary we may assume that w lies in the interior of σ , hence $0 < \alpha_{ij}$. Since $f(w) \in N_\tau$, it follows that $\sum_j \alpha_{ij} \in \mathbb{Z}$ for all i . In particular, if $J_{i_0} = 1$ for some i_0 then $\alpha_{i_0 1} = 0$, and w lies in a face of σ . So we may assume that $J_i > 1$ for all i . Since the relative dimension of f is $\sum_i (J_i - 1)$, we have to consider all possible decompositions $\sum_i (J_i - 1) \leq 3$, where $J_i > 1$ for all i .

The cases when the relative dimension of f is 0 or 1 are trivial and left to the reader.

If the relative dimension of f is 2, then either $J_1 = 3$, or $J_1 = J_2 = 2$. In the first case, we have that $\langle v_{11}, v_{12}, v_{13} \rangle$ is singular, contradicting the semistability of $f|_{\Delta_{X,1}}$. In the second case, $\alpha_{11} + \alpha_{12}, \alpha_{21} + \alpha_{22} \in \mathbb{Z}$ and $0 < \alpha_{ij} < 1$ imply that $\alpha_{11} + \alpha_{12} = \alpha_{21} + \alpha_{22} = 1$. Hence $f(w) = u_1 + u_2$ is a barycenter.

In relative dimension 3, either $J_1 = 4$, or $J_1 = 3, J_2 = 2$, or $J_1 = J_2 = J_3 = 2$. In the first case, we get a contradiction with the semistability of $f|_{\Delta_{X,1}}$; the third case gives $\alpha_{11} + \alpha_{12} = \alpha_{21} + \alpha_{22} = \alpha_{31} + \alpha_{32} = 1$ as for relative dimension 2. In the second case either $\alpha_{11} + \alpha_{12} + \alpha_{13} = \alpha_{21} + \alpha_{22} = 1$ and w maps to a barycenter, or $\alpha_{11} + \alpha_{12} + \alpha_{13} = 2, \alpha_{21} + \alpha_{22} = 1$ and $(\sum v_{ij}) - w$ maps to a barycenter. $\square$

REFERENCES

- [N-K] D. Abramovich and K. Karu, *Weak semistable reduction in characteristic 0*, preprint. [alg-geom/9707012](https://arxiv.org/abs/alg-geom/9707012).
- [F] W. Fulton, *Introduction to Toric Varieties*, Princeton University Press, 1993.
- [KKMS] G. Kempf, F. Knudsen, D. Mumford and B. Saint-Donat, *Toroidal Embeddings I*, Springer, LNM 339, 1973.

DEPARTMENT OF MATHEMATICS, BOSTON UNIVERSITY, 111 CUMMINGTON, BOSTON, MA 02215, USA

E-mail address: `kllkr@math.bu.edu`